



## Quantitative pharmacologic MRI: Mapping the cerebral blood volume response to cocaine in dopamine transporter knockout mice

Teodora-Adriana Perles-Barbacaru<sup>a</sup>, Daniel Procissi<sup>a</sup>, Andrey V. Demyanenko<sup>a</sup>, F. Scott Hall<sup>b</sup>, George R. Uhl<sup>b</sup>, Russell E. Jacobs<sup>a,\*</sup>

<sup>a</sup> Biological Imaging Center, Beckman Institute, California Institute of Technology, Caltech M/C 139-74, 1200 E. California Blvd, Pasadena, CA 91125, USA

<sup>b</sup> Molecular Neurobiology Branch, National Institute on Drug Abuse, Intramural Research Program, Baltimore, MD 21224, USA

### ARTICLE INFO

#### Article history:

Received 17 May 2010

Revised 15 December 2010

Accepted 17 December 2010

Available online 23 December 2010

### ABSTRACT

The use of pharmacologic MRI (phMRI) in mouse models of brain disorders allows noninvasive *in vivo* assessment of drug-modulated local cerebral blood volume changes ( $\Delta$ CBV) as one correlate of neuronal and neurovascular activities. In this report, we employed CBV-weighted phMRI to compare cocaine-modulated neuronal activity in dopamine transporter (DAT) knockout (KO) and wild-type mice. Cocaine acts to block the dopamine, norepinephrine, and serotonin transporters (DAT, NET, and SERT) that clear their respective neurotransmitters from the synapses, helping to terminate cognate neurotransmission. Cocaine consistently reduced CBV, with a similar pattern of regional  $\Delta$ CBV in brain structures involved in mediating reward in both DAT genotypes. The largest effects ( $-20\%$  to  $-30\%$   $\Delta$ CBV) were seen in the nucleus accumbens and several cortical regions. Decreasing response amplitudes to cocaine were noted in more posterior components of the cortico-mesolimbic circuit. DAT KO mice had significantly attenuated  $\Delta$ CBV amplitudes, shortened times to peak response, and reduced response duration in most regions. This study demonstrates that DAT knockout does not abolish the phMRI responses to cocaine, suggesting that adaptations to loss of DAT and/or retained cocaine activity in other monoamine neurotransmitter systems underlie these responses in DAT KO mice.

© 2010 Elsevier Inc. All rights reserved.

### Introduction

Assessment of the effects of psychotropic substances requires an understanding of their sites, mechanisms, and time courses of action. Functional neuroimaging techniques have made it possible to study drug-induced changes in brain activity *in vivo* (Leslie and James, 2000). Pharmacological magnetic resonance imaging (phMRI), a variant of functional MRI, is well adapted to monitor drug action in the brain dynamically and noninvasively and to map the brain systems and networks targeted by psychoactive drugs (Chen et al., 2001). A number of important classes of medications used in the treatment of psychiatric disorders and many drugs of abuse target plasma membrane reuptake transporters for the monoamine neurotransmitters dopamine, norepinephrine, and serotonin (DAT, NET, and SERT, respectively) inhibiting the reuptake of the released neurotransmitters (Gainetdinov and Caron, 2003).

Dopamine systems, including the cortico-mesolimbic system, are crucial for normal reward-seeking behavior and are likely to make major contributions to the development of addiction (Girault and Greengard, 2004). Cocaine is a prototypical psychomotor stimulant with powerful mood altering properties (Johanson and Fischman, 1989; Uhl et al.,

2002). It has dramatic effects on the reward pathway through interaction with the monoamine transporters (Gainetdinov and Caron, 2003). Moreover, chronic exposure has a multitude of effects in the brain, including modulation of signaling cascades, neurotrophic factors, monoamine receptors, and neuroplasticity (El-Ghundi et al., 2007; Thomas et al., 2008). Contrary to initial expectations, DAT knockout (KO) mice display evidence for cocaine reinforcement in conditioned place preference and even in some self-administration paradigms (Rocha et al., 1998; Sora et al., 1998). However, in one line of DAT KO mice, individual mice differ somewhat in their cocaine self-administration behavior. Most do not self-administer. Those that do self-administer fail to continue responding when challenged with more demanding schedules of reinforcement (Thomsen et al., 2009). This may be due to disruption of aspects of incentive motivation mediated by dopamine, since DAT KO mice do not exhibit cocaine conditioned locomotion (Hall et al., 2009). That cocaine is rewarding at all in DAT KO mice may be, in part, a consequence of neuroadaptations in the SERT and NET systems, consistent with the finding that SERT and NET blockers have reinforcing effects in DAT KO mice that are not observed in wild-type mice (Hall et al., 2002). Such alterations might also involve changes in the aversive effects of cocaine (Uhl et al., 2002). Overall, these data suggest that although aspects of cocaine reward are maintained, at least to a certain degree, in DAT KO mice, the mechanisms underlying cocaine reward differ from those in wild-type mice.

\* Corresponding author. Fax: +1 626 449 5163.

E-mail address: [rjacobs@caltech.edu](mailto:rjacobs@caltech.edu) (R.E. Jacobs).

phMRI measures the hemodynamic responses to a pharmacological challenge and thus probes the activity of neuronal pathways targeted by the drug based on the neurovascular consequences of changed neuronal (and vascular) activities (Jenkins et al., 2004; Mandeville et al., 1998; Schwarz et al., 2007a). This technique has the potential for examining the underlying changes in regional brain activity in mouse models of disorders that display links to altered neurotransmitter activity, including attention deficit hyperactivity disorder, narcolepsy, obesity, and drug addiction. However, small brain size, rapid metabolic rates, susceptibility artifacts, and the need for anesthesia render functional MRI challenging in mice. Small animal imaging is carried out at high magnetic field strengths (>7 T) to maximize signal intensity. In phMRI, the sensitivity to functional changes is enhanced with exogenous magnetic resonance contrast agents such as ultrasmall superparamagnetic iron oxide particles (USPIO) that are retained in the blood pool and exhibit long circulation times (Weissleder et al., 1990). The paramagnetic signals generated by such superparamagnetic particles are so large that most of the MRI signal is determined by the cerebral blood volume (CBV), overwhelming blood oxygen level-dependent (BOLD), and blood flow effects. Most current studies using this technique have used rats (Mandeville et al., 2004, 1998; Schwarz et al., 2007b) or nonhuman primates (Jenkins et al., 2004). A few research teams have successfully performed CBV-weighted phMRI in mice (Luo et al., 2008; Mueggler et al., 2003; Wu et al., 2004), although low signal-to-noise ratio (SNR) and sensitivity have precluded a broad range of investigations using genetically engineered mouse models.

In the present report, we have examined, for the first time, effects of DAT gene deletion on neurovascular responses to an acute cocaine challenge. We have used a CBV-weighted phMRI technique in mice and achieved good sensitivity to the CBV change with time resolution of 1 min and spatial resolution of <0.2 mm. This methodology has already proved useful in larger rodent and primate systems (Chin et al., 2006; Jenkins et al., 2004; Mandeville et al., 2005; Marota et al., 2000; Sanchez-Pernaute et al., 2007; Schwarz et al., 2003). Adapting this approach for mouse brain imaging should provide a valuable means for testing hypotheses in a variety of mouse model systems.

## Materials and methods

CBV responses to cocaine were studied in 6 DAT KO mice and 6 wild-type control mice consisting of 4 wild-type littermates and 2 C57BL/6J female mice (The Jackson Laboratory, Bar Harbor, ME). The DAT KO mice used in these experiments have been described previously (Sora et al., 1998). Pilot experiments on isoflurane anesthetized mice of different strains and ages were carried out to establish a suitable cocaine dose. In bench top experiments that did not involve MRI, the electrocardiogram, respiration, and expired CO<sub>2</sub> (MicroCapStar CO<sub>2</sub> Monitor, IITC Life Science Inc., CA) were continuously recorded before and after cocaine injection. Criteria for the choice of the cocaine dose were (a) no arousal from anesthesia and (b) stable physiologic parameters after injection. Behavioral changes have been reported starting at a cocaine dose of 5 mg/kg intraperitoneal (IP) in conscious mice (Jamshidi et al., 2004), while cocaine-induced seizures start to appear at a dose of 70 mg/kg IP. Doses of 20 to 40 mg/kg IP increase striatal and nucleus accumbens dopamine levels 3-fold in wild-type mice (Mateo et al., 2004; Rocha et al., 1998). In this work, cocaine hydrochloride (Sigma-Aldrich, MO) was dissolved in normal saline solution (5 mg/ml) for IP injection at a dose of 30 mg/kg.

P904 (Guerbet Research, Aulnay-Sous-Bois, France) is a research prototype USPIO with a hydrodynamic diameter of 25 to 30 nm and a glucose derivative coating designed for macrophage and vascular imaging applications (Sigovan et al., 2009). It was

provided as a 500 mM (≈28 mg/ml) solution and was diluted to 6 mg/ml in sterile normal saline solution for IV injection. A volume of ≈4.2 μl/g body weight was injected IV resulting in a dose of 25 mg/kg.

All experiments were performed in accordance with protocols approved by the Institutional Animal Care and Use Committee of the California Institute of Technology.

## *In vivo* phMRI

*In vivo* MRI experiments were performed in a horizontal bore Bruker Biospec/Avance 7T/30 cm small animal MR system using a custom made 12-rung linear birdcage RF coil (length 45 mm, inner diameter 26 mm) for transmission and reception. The mice were positioned prone in a plastic cradle with an integral head mask providing 1.5 to 1.7% isoflurane in N<sub>2</sub>/O<sub>2</sub> 70%/30% and incorporated ECG electrodes for the front paws and tail. Rectal temperature (Opsens OTG-M, Canada) and respiration (BIOPAC Systems, CA) were monitored continuously. Simultaneous recording of the electrocardiogram during gradient echo imaging was not always reliable, but heart rates at the start and end of acquisition were comparable. Body temperature was maintained between 36 and 38 °C using heated air passed through the RF coil and a cylindrical plastic extension that covered the mouse body, preventing rapid loss of the warm air. The isoflurane level was adjusted to achieve respiratory rates of 80 to 120 breaths per minute and was not changed during image acquisition. The mice were fitted with tail vein catheters for USPIO injection and intraperitoneal catheters for drug injection. Both catheters had extension lines with total volumes of 200 μl and were preloaded with USPIO and drug in normal saline solution before positioning the mouse in the magnet.

An autoshimming routine resulted in a <sup>1</sup>H linewidth of <0.13 ppm (FWHM) in a cube of 4 × 4 × 4 mm<sup>3</sup> brain tissue. After obtaining stable physiologic conditions, relative CBV maps were obtained based on the intravascular susceptibility effect of the USPIO (Berry et al., 1996; Mandeville et al., 1998).

For each brain, 14 contiguous axial 0.75-mm-thick slices, with a field of view of 19 × 19 mm<sup>2</sup>, were acquired from the olfactory bulb to the cerebellum using a 2D multiple gradient echo sequence with TR = 600 ms, flip angle = 35°, 100 × 100 matrix, spectral line width 60 kHz, three echoes at TE = 2.5, 6.0 and 9.5 ms, and an acquisition time of 1 min per repetition.

Fig. 1 illustrates the phMRI protocol: R<sub>2</sub>\*-weighted images were acquired for 30 min prior to contrast agent administration followed by 30 post-contrast baseline acquisitions. After the pharmacological challenge, image acquisition continued for 90 min. The USPIO and cocaine were injected by flushing the IV/IP catheter with 250 μl saline solution over a duration of 1 min while the mouse was positioned inside the scanner.


## MRI data analysis

Pre-contrast and post-contrast images were automatically co-registered with a 2D rigid body (3 parameters) model using Automated Image Registration (AIR) software (Woods et al., 1998). Image analysis was performed with ImageJ (Abramoff et al., 2004). No spatial or temporal smoothing was applied to the data.

The relative signal decrease upon contrast agent administration is

$$1 - \frac{S_{\text{postCA}}}{S_{\text{preCA}}}, \quad (1)$$

where  $S_{\text{preCA}}$  is the mean signal intensity prior to contrast agent (CA) injection and  $S_{\text{postCA}}$  is the signal intensity averaged over the first five acquisitions after contrast agent injection (Fig. 1). The signal decrease depends on the relaxivity of the contrast agent and its initial tissue



**Fig. 1.** Experimental protocol: in pHMRI experiments gradient echo images were acquired continuously with a time resolution of 1 min. The baseline intervals before and after P904 injection were 30 min each.  $S_{preCA}$  is the constant baseline signal before contrast agent injection;  $S_{postCA}$  is the average of five consecutive signals acquired immediately after contrast agent injection, and  $S(t)$  is the time dependent CBV-weighted signal after contrast agent injection. The acquisitions in the interval between contrast agent and cocaine injection and the acquisitions at >60 min after cocaine were used to detrend the signal and to derive the contrast agent washout rate.

concentration. These factors determine the transverse relaxation rate change

$$\Delta R_2^* = -\frac{1}{TE} \ln\left(\frac{S_{postCA}}{S_{preCA}}\right). \quad (2)$$

where TE is the echo time.

Prior to quantitative analysis, the signal intensity after contrast agent injection,  $S(t)$ , was detrended (Schwarz et al., 2003) to eliminate the effect of contrast agent washout by fitting two time intervals of  $S(t)$  prior to and >60 min after pharmacological challenge (Fig. 1) to a two-parameter model (Perles-Barbacaru et al., 2010). Assuming that the  $T_2$ -weighted signal is a simple exponential function ( $S_0(t) \propto \exp[-TE R_2^*(t)]$ ), we find that:

$$S_0(t) = S_{preCA} \exp\left(-TE \cdot \kappa \cdot \exp\left(\frac{t}{\tau}\right)\right) \quad (3)$$

where  $S_{preCA}$  is defined as above. The fitting parameter  $\kappa$  provides an estimate of the initial  $\Delta R_2^*$  following USPIO injection (Eq. (2)), while  $\tau^{-1}$  is the apparent washout rate of the contrast agent from the tissue. In this model, we assume negligible  $T_1$  effects, an instantaneous USPIO bolus and mono-exponential USPIO washout in the plasma, since USPIO is confined to the intravascular compartment.

The relative CBV change ( $\Delta CBV$ ) is calculated for each image voxel according to Mandeville et al. (1998) by assuming a linear relationship between CBV and  $\Delta R_2^*$ :

$$\Delta CBV(t) = \frac{\ln(S_0(t)/S(t))}{\ln\left(\frac{S_{preCA}}{S_0(t)}\right)} \quad (4)$$

where  $S_0(t)$  represents the fit to the signal time course after contrast agent injection under resting conditions with no drug challenge (Eq. (3)).  $S(t)$  refers to the measured signal after contrast agent injection. We analyzed the  $\Delta CBV$  using the signal from the 3rd echo at TE = 9.5 ms, while the ROIs were drawn on the images of the first two echoes.

Quantitative analyses of the parametric maps from Eq. (4) describing the CBV change upon cocaine injection were performed by examining several cerebral ROIs forming the drug reward circuitry. We analyzed ROIs in the prefrontal, frontal, and frontoparietal cortex (CTX); nucleus accumbens (ACB); caudate putamen (CP); amygdala (AMY); hippocampal formation (HPF); thalamic nuclei (TH); hypothalamus (HY); midbrain (MB); ventral tegmental area (VTA); substantia nigra (SN); pons (P); medulla (MY); and cerebellum (CBX). To rule out confounding systemic vascular changes associated with the pharmacological stimulus and independent of neuronal activity, we also analyzed the signal change in an extra-cerebral ROI located in the masseter or temporalis muscle.

To describe the regional time course of the  $\Delta CBV$  response, we used a simplified gamma variate model.

$$\Delta CBV(t) = \Delta CBV_{peak} \left(\frac{t}{T_{peak}}\right)^\alpha \exp\left(\alpha \left(1 - \frac{t}{T_{peak}}\right)\right) \quad (5)$$


The three model parameters ( $\Delta CBV_{peak}$  = amplitude,  $T_{peak}$  = time to peak, and  $\alpha$  = shape factor) were compared between DAT KO and wild-type control groups using unpaired  $T$ -tests. In particular, for each ROI noted above, the gamma variate model (Eq. (5)) was fitted to the data of one animal at a time and to the combined data from all animals in one group (examples for ACB and VTA can be seen in Fig. 3). An  $F$ -test was used to compare the relative difference in the sum of squares of the residuals from each fit and the relative difference in the degrees of freedom. To test for significant differences in each model parameter, an  $F$ -test was applied to the results of nonlinear regression with independent and shared parameter of interest.  $T$ -tests for each model parameter confirmed the statistically significant differences detected by the  $F$ -test. To test whether the entire response curves were different, the  $F$ -test was performed on the results of the nonlinear regression to the separate data and to the combined data of both groups.  $P < 0.01$  was considered statistically significant. Curve fitting and statistical tests were performed using GraphPad Prism version 4.03 for Windows (GraphPad Software, La Jolla, CA).

## Results

During and within 1 min after intravenous injection of P904, respiratory amplitudes and rates were increased. Pilot experiments showed that cocaine doses up to 1.5 mg/kg IV or 30 mg/kg IP did not alter these physiologic parameters for more than 3 min. We administered cocaine intraperitoneally during MRI experiments to avoid sudden hypervolemia and to avoid the tail and body motion that often accompanied intravenous cocaine administration. An intraperitoneal dose of 30 mg/kg cocaine transiently reduced and then increased the respiratory amplitude for 1 to 3 min.

Analysis of the signal from a ROI in the dorsal striatum (area  $3.2 \pm 0.2 \text{ mm}^2$ ) acquired at TE = 9.5 ms showed that a P904 dose of 25 mg/kg yielded a relative signal decrease (Eq. (1)) of  $0.58 \pm 0.02$  during the 5 min following contrast agent injection with an apparent tissue half-life ( $\tau \cdot \ln 2$ ) of  $156 \pm 19 \text{ min}$  ( $N = 12$ , average  $\pm$  standard error of the mean). Although the apparent tissue half-life computed according to Equation (3) is region dependent, it is greater than 2 hours in all regions. Thus, the detrending model in Eq. (3) takes reasonable account of the slowly decreasing P904 concentration.

Saline injection produced no measurable effect on CBV (data not shown). Fig. 2 shows that a spatially and temporally dependent CBV change was observed in cerebral tissue of mice treated with cocaine, while there was no measurable effect on extra-cerebral vasculature. Cocaine induced a marked CBV decrease (Fig. 2) that lasted between 40 and 60 min. The ventral striatum/nucleus accumbens exhibited the strongest response to cocaine.




**Fig. 2.** Effect of IP cocaine on the signal time course in cerebral (nucleus accumbens, open symbols) and in extra-cerebral tissue (muscle, closed symbols). Injection of 25 mg/kg P904 IV at  $t = -30$  min (left arrow) yields a 62.5% signal drop in nucleus accumbens ( $4.06 \text{ mm}^3$ ) and 69.1% signal drop in muscle ( $9.58 \text{ mm}^3$ ) at  $TE = 9.5$  ms. The second arrow indicates the IP injection of cocaine (30 mg/kg), which leads to a signal increase in the cerebral tissue corresponding to a CBV decrease. The error bars show the standard deviation of the signal within the ROI in a single mouse. This is a typical signal time course prior to detrending.

The time course of the  $\Delta\text{CBV}$  response to cocaine in two ROIs within the mesolimbic pathway is shown in Fig. 3 for DAT KO and wild-type mice. The time courses in all examined ROIs, except thalamic nuclei, differed significantly between DAT genotypes. Fig. 4 shows that the DAT KO mice display an attenuated  $\Delta\text{CBV}$  response in all regions except the hypothalamus, thalamus, pons, and medulla oblongata. Also, in the knockout mice, peak responses were reached at earlier time points for most brain structures and the total duration of the hemodynamic response to cocaine was shorter than in wild-type mice (Figs. 3 and 4).

## Discussion

### Drug-induced CBV changes

Species differences in cerebral responses to cocaine have been reported previously (Lyons et al., 1996; Zocchi et al., 2001) but



**Fig. 3.**  $\Delta\text{CBV}$  response in two cerebral structures (nucleus accumbens and ventral tegmental area) to an IP injection of 30 mg/kg cocaine at  $t = 0$ . The error bars represent SEM;  $N = 6$  DAT KO and  $N = 6$  wild-type mice. The continuous black lines are the best gamma variate fit with the 95% confidence interval shown as dashed gray lines.

remain unexplained. In this study, a negative  $\Delta\text{CBV}$  response to cocaine was observed in mice. In rats,  $\Delta\text{CBV}$  increases after cocaine challenge have been observed with MRI (Ceolin et al., 2007; Marota et al., 2000; Schwarz et al., 2003, 2004), while cocaine-induced  $\Delta\text{CBV}$  decreases have been observed in humans (Kaufman et al., 1998). Mechanisms for the negative CBV response may be related to reduced cerebrovascular feedback and/or decreased neuronal activation (Mandeville et al., 2005; Peoples et al., 1998) accompanied by a decreased cerebral glucose metabolism (Lyons et al., 1996; Thanos et al., 2008). Cerebral microvascular constriction has also been observed in rhesus monkey brain slices upon micro-application of dopamine (Krimer et al., 1998). Direct activation of D1 dopaminergic receptors on arterioles results in vasodilation (Choi et al., 2006), although the sign of the CBV response depends on both the effective concentration of dopamine and the relative regional densities of receptors in the D1/D5 and D2/D3 and possibly even D4 receptor families (Choi et al., 2006). Cocaine-induced blockade of norepinephrine and serotonin reuptake are also likely to influence the overall direction of cardiovascular changes in both the wild type, perhaps especially in the DAT KO mice.


### Confounding vascular effects

Vascular effects not related to neuroactivity also have to be considered when interpreting pHMRI results. Confounding systemic effects of cocaine have been difficult to exclude (Gozzi et al., 2007; Luo et al., 2009). A limitation of the current study is that the arterial blood pressure could not be monitored during scanning. However, the transient effect ( $\sim 5$  min) of this cocaine dose on the systemic blood pressure does not seem to provide a large determinant for the sustained ( $\sim 1$  h) hemodynamic response in the brain (Du et al., 2009, 2006; Luo et al., 2003; Schmidt et al., 2006). Moreover, any large peripheral blood pressure change following cocaine injection would have resulted in a signal change in muscle tissue. Our findings suggest that the measured CBV change in the current study is largely independent of systemic effects of the drug.

There is evidence that the CBV response may depend on the type of anesthetic used in the experiment (Du et al., 2009; Luo et al., 2009). In previous studies, which used different anesthetics, the anesthetic used may have been confounded with apparent species differences. In a study by Du et al. (2006), a CBV decrease of  $-4\%$  to  $-8\%$  was observed with optical techniques in the cortex of rats under isoflurane anesthesia after an intravenous dose of 1 mg/kg cocaine whether or not a peripheral blood pressure decrease during the first minutes was prevented by phenylephrine. Consistent anesthesia conditions were employed throughout the present work. Comparisons among the different genotypes are thus credible, regardless of whether the nature of differences in CBV responses between previous studies is the result of true species differences or procedural differences (e.g., the choice of anesthetic). Nevertheless, care must be taken in comparisons with conscious animals or with results from studies that use other anesthesia regimens.

### Reward circuitry and monoamine transporter knockout mice

The cortico-mesolimbic dopaminergic circuit is thought to mediate many of the rewarding effects of many drugs of abuse, including cocaine, which prolongs dopaminergic transmission. However, cocaine also elevates extracellular serotonin and norepinephrine levels (Li et al., 1996), which may account for its pronounced rewarding effects compared to more selective agents. Rodents, in general, and monoamine transporter knockout mice, in particular, have been used in efforts to identify the mechanisms of action of cocaine and other drugs of abuse. A study by Schwarz and colleagues in rats (Schwarz et al., 2004) has shown that the  $\Delta\text{CBV}$  time course is not correlated with the local dopamine concentration or with the local cocaine



**Fig. 4.** Quantitative analysis of the regional CBV change after cocaine injection. Peak amplitude and the time to peak of the gamma variate fit to the signal time course are shown for specific brain regions. The error bars indicate the standard error of the mean ( $N = 6$  per group). In all ROIs, there is a statistically significant difference ( $P < 0.01$ ) between DAT KO and wild-type mice in at least one of the three fitting parameters (the shape factor  $\alpha$  is not shown). The overall time course of the  $\Delta$ CBV response is statistically different for KO and control mice in all ROIs except the thalamus. Typical ROI outlines are shown on two coronal brain slices: prefrontal, frontal, and frontoparietal cortex (CTX); nucleus accumbens (ACB); caudate putamen (CP); amygdala (AMY); hippocampal formation (HPF); thalamic nuclei (TH); hypothalamus (HY); midbrain (MB); ventral tegmental area (VTA); substantia nigra (SN); pons (P); medulla (MY); and cerebellum (CBX).

concentration in all brain regions. A more complex mechanism for the  $\Delta$ CBV response involving other neurotransmitters and downstream consequences of the effects produced by cocaine is supported by the fact that, in our study, a straightforward pharmacokinetic model did not fit the  $\Delta$ CBV time course. The  $\Delta$ CBV response was, therefore, described by a gamma variate model that allowed for a gradual onset of the  $\Delta$ CBV response.

In this study, similar patterns of spatial activation were observed in both DAT KO and wild-type mice. The largest negative responses were observed in the nucleus accumbens, frontal and parietal cortex, dorsal striatum, and amygdala (in decreasing order). These structures have substantial dopaminergic innervation and constitute a substantial portion of the brain reward circuitry that mediates the rewarding effects of drugs of abuse (Koob and Volkow, 2010). The time to peak response was 12–18 min in all structures examined, except the cerebellum where it was 20–28 min. Quantitative regional analysis revealed a significant decrease in amplitude and a slight decrease in duration of the hemodynamic response in DAT KO mice compared to wild-type mice. The  $\Delta$ CBV response to cocaine in these regions was attenuated but not entirely abolished in DAT KO mice. This is consistent with the maintenance of some reinforcing effects of cocaine in DAT KO mice at the cocaine dose used here as indicated by conditioned place preference (Sora et al., 2001, 1998) and cocaine self-administration (Rocha et al., 1998) studies. However, the reduced  $\Delta$ CBV response in DAT KO mice may also be considered to be consistent with the weaker reinforcing effects that have been demonstrated in a more recent self-administration study (Thomsen et al., 2009). More importantly, it may indicate a difference in the underlying mechanism of cocaine effects in DAT KO and wild-type mice. Cocaine continues to produce a conditioned place preference in DAT KO mice, but DAT KO mice also exhibit novel rewarding effects of selective SERT and NET blockers (Hall et al., 2002). The rewarding effects of these drugs may be due to consequences of SERT and NET blockade in the VTA/SN. Along with its rewarding effects, cocaine continues to produce increases in extracellular dopamine in the striatum in this strain of DAT KO mice (Shen et al., 2004). This might be due to cocaine blockade of SERT, a hypothesis consistent with the elimination of cocaine conditioned place preference in DAT/SERT double KO mice (Sora et al., 2001). Furthermore, although fluoxetine, a selective serotonin reuptake inhibitor, has no effect in wild-type mice, it is rewarding (Hall et al., 2002) and increases striatal dopamine levels in DAT KO mice (Shen et al., 2004). The locus of this effect may

be the VTA/SN, as local infusions of cocaine in either the dorsal or ventral striatum do not increase extracellular dopamine levels in DAT KO mice (Mateo et al., 2004; Shen et al., 2004), but local injections of cocaine or fluoxetine into the VTA do increase extracellular dopamine levels in the nucleus accumbens of DAT KO, but not wild-type, mice (Mateo et al., 2004). In the present study, the greatest differences in  $\Delta$ CBV between KO and wild-type mice were observed in the VTA/SN. These results are consistent with the hypothesis presented above that the mechanism of cocaine-induced increases in striatal dopamine in DAT KO mice involve at least some cocaine actions in the VTA/SN rather than dopaminergic terminal fields.

DAT KO responses to cocaine are less negative than those of wild-type animals in all brain regions except the pons, hypothalamus, and medulla oblongata (and not statistically different in the thalamus). These are all regions that are not involved in drug reward. They display only modest densities of dopaminergic innervation. Thanos et al. (2008) used PET measured  $^{18}$ F-fluorodeoxyglucose ( $^{18}$ FDG) uptake in DAT KO and wild-type mice to assay metabolic activity. They found an overall decrease in baseline  $^{18}$ FDG uptake in DAT KO versus wild-type mice, and increased uptake after cocaine challenge in DAT KO mice only in the thalamus and cerebellum. In that PET study  $^{18}$ FDG was introduced 30 min after cocaine injection with an 80-min acquisition time. Thus, the  $^{18}$ FDG uptake reflects cocaine effects well after the CBV changes observed here have returned to baseline. In other studies with rats and primates using intravenous rather than IP injection of drugs (Jenkins et al., 2004; Marota et al., 2000), both time to maximum change in CBV and return to baseline are significantly reduced from values observed here due to different routes of administration. Regardless of how cocaine is administered,  $\Delta$ CBV changes measure a fast response of the reward circuit to cocaine challenge, while  $^{18}$ FDG uptake evaluates longer term responses, averaged over 80 min, that may reflect adaptations to the initial cocaine effect. Taken together, the PET and pHMRI results imply that cocaine impacts the cortico-mesolimbic reward circuit in the first minutes after challenge with the most impact in the more dopaminergically innervated anterior structures. Cocaine subsequently exerts a more sustained influence on the more “limbic” parts of the circuit.

## Conclusion

This study demonstrates the applicability of the CBV-weighted pHMRI technique to mapping hemodynamic effects of centrally acting

drugs in mouse models of disease. The technique is minimally invasive and provides good functional sensitivity. We also note that, as with other functional MRI techniques, the mechanisms underlying the CBV changes may not be exclusively specific to neuronal activation. The negative  $\Delta$ CBV response to cocaine challenge is rapid (~15 min) throughout the reward circuit and attenuated in DAT KO mice. Since knockout of DAT has a small but significant effect on the immediate response to cocaine in brain regions thought to be involved in cocaine reward, knockout of DAT appears to produce circuit-wide changes, not just changes in dopaminergic neurons. The largest responses to cocaine (–20% to –30%  $\Delta$ CBV) are seen in the cortex and nucleus accumbens. There are decreasing responses in more posterior parts of the reward circuit (CP>AMY>VTA~SN). Nevertheless, some of the greatest differences between DAT KO and wild-type mice were observed in midbrain structures, VTA and SN, which may indicate a different mechanism of action of cocaine in DAT KO mice. The fact that there are much more modest differences in the nucleus accumbens and dorsal striatum between DAT KO and wild-type mice is consistent with the retention of some cocaine reward, under at least some conditions (Rocha et al., 1998; Sora et al., 2001, 1998; Thomsen et al., 2009). Thus these results emphasize the ultimate importance of the nucleus accumbens at early as well as later stages of drug exposure (Koob and Volkow, 2010).

## Acknowledgments

The authors thank Hargun Sohi and Thomas Ng for their technical assistance, Davit Janvelyan for helping with the AIR software, Guerbet Research (Aulnay-Sous-Bois, France) for providing the P904 contrast agent. This project was funded in part by the Beckman Institute, NIDA R01DA18184, and NCRRT U24 RR021760 Mouse BIRN and by funding from the National Institute on Drug Abuse, Intramural Research Program (GRU, FSH).

## References

- Abramoff, M.D., Magelhaes, P.J., Ram, S.J., 2004. Image processing with Image. *J. Biophotonics Int.* 11, 36–42.
- Berry, I., Benderbous, S., Ranjeva, J.P., Gracia-Meavilla, D., Manelfe, C., Le Bihan, D., 1996. Contribution of Sinerem used as blood-pool contrast agent: detection of cerebral blood volume changes during apnea in the rabbit. *Magn. Reson. Med.* 36, 415–419.
- Ceolin, L., Schwarz, A.J., Gozzi, A., Reese, T., Bifone, A., 2007. Effects of cocaine on blood flow and oxygen metabolism in the rat brain: implications for pHfMRI. *Magn. Reson. Imaging* 25, 795–800.
- Chen, Y.C., Mandeville, J.B., Nguyen, T.V., Talele, A., Cavagna, F., Jenkins, B.G., 2001. Improved mapping of pharmacologically induced neuronal activation using the IRON technique with superparamagnetic blood pool agents. *J. Magn. Reson. Imaging* 14, 517–524.
- Chin, C.L., Fox, G.B., Hradil, V.P., Osinski, M.A., McGaraughty, S.P., Skoubis, P.D., Cox, B.F., Luo, Y., 2006. Pharmacological MRI in awake rats reveals neural activity in area postrema and nucleus tractus solitarius: relevance as a potential biomarker for detecting drug-induced emesis. *Neuroimage* 33, 1152–1160.
- Choi, J.K., Chen, Y.I., Hamel, E., Jenkins, B.G., 2006. Brain hemodynamic changes mediated by dopamine receptors: role of the cerebral microvasculature in dopamine-mediated neurovascular coupling. *Neuroimage* 30, 700–712.
- Du, C., Tully, M., Volkow, N.D., Schiffer, W.K., Yu, M., Luo, Z., Koretsky, A.P., Benveniste, H., 2009. Differential effects of anesthetics on cocaine's pharmacokinetic and pharmacodynamic effects in brain. *Eur. J. Neurosci.* 30, 1565–1575.
- Du, C., Yu, M., Volkow, N.D., Koretsky, A.P., Fowler, J.S., Benveniste, H., 2006. Cocaine increases the intracellular calcium concentration in brain independently of its cerebrovascular effects. *J. Neurosci.* 26, 11522–11531.
- El-Ghundi, M., O'Dowd, B.F., George, S.R., 2007. Insights into the role of dopamine receptor systems in learning and memory. *Rev. Neurosci.* 18, 37–66.
- Gainetdinov, R.R., Caron, M.G., 2003. Monoamine transporters: from genes to behavior. *Annu. Rev. Pharmacol. Toxicol.* 43, 261–284.
- Girault, J.A., Greengard, P., 2004. The neurobiology of dopamine signaling. *Arch. Neurol.* 61, 641–644.
- Gozzi, A., Ceolin, L., Schwarz, A., Reese, T., Bertani, S., Crestan, V., Bifone, A., 2007. A multimodality investigation of cerebral hemodynamics and autoregulation in pharmacological MRI. *Magn. Reson. Imaging* 25, 826–833.
- Hall, F.S., Li, X.F., Randall-Thompson, J., Sora, I., Murphy, D.L., Lesch, K.P., Caron, M., Uhl, G.R., 2009. Cocaine-conditioned locomotion in dopamine transporter, norepinephrine transporter and 5-HT transporter knockout mice. *Neuroscience* 162, 870–880.
- Hall, F.S., Li, X.F., Sora, I., Xu, F., Caron, M., Lesch, K.P., Murphy, D.L., Uhl, G.R., 2002. Cocaine mechanisms: enhanced cocaine, fluoxetine and nisoxetine place preferences following monoamine transporter deletions. *Neuroscience* 115, 153–161.
- Jamshidi, H.R., Rezayat, M., Zarrindast, M.R., 2004. Effect of ampin on tolerance to cocaine-induced locomotor activity in mice. *Acta Med. Iran.* 42, 78–82.
- Jenkins, B.G., Sanchez-Pernaute, R., Brownell, A.-L., Chen, Y.-C.I., Isacson, O., 2004. Mapping dopamine function in primates using pharmacologic magnetic resonance imaging. *J. Neurosci.* 24, 9553–9560.
- Johanson, C.E., Fischman, M.W., 1989. The pharmacology of cocaine related to its abuse. *Pharmacol. Rev.* 41, 3–52.
- Kaufman, M.J., Levin, J.M., Maas, L.C., Rose, S.L., Lukas, S.E., Mendelson, J.H., Cohen, B.M., Renshaw, P.F., 1998. Cocaine decreases relative cerebral blood volume in humans: a dynamic susceptibility contrast magnetic resonance imaging study. *Psychopharmacology* 138, 76–81.
- Koob, G.F., Volkow, N.D., 2010. Neurocircuitry of addiction. *Neuropsychopharmacology* 35, 217–238.
- Krimer, L.S., Muly III, E.C., Williams, G.V., Goldman-Rakic, P.S., 1998. Dopaminergic regulation of cerebral cortical microcirculation. *Nat. Neurosci.* 1, 286–289.
- Leslie, R.A., James, M.F., 2000. Pharmacological magnetic resonance imaging: a new application for functional MRI. *Trends Pharmacol. Sci.* 21, 314–318.
- Li, M.Y., Yan, Q.S., Coffey, L.L., Reith, M.E.A., 1996. Extracellular dopamine, norepinephrine, and serotonin in the nucleus accumbens of freely moving rats during intracerebral dialysis with cocaine and other monoamine uptake blockers. *J. Neurochem.* 66, 559–568.
- Luo, F., Seifert, T.R., Edalji, R., Loebbert, R.W., Hradil, V.P., Harlan, J., Schmidt, M., Nimmrich, V., Cox, B.F., Fox, G.B., 2008. Non-invasive characterization of beta-amyloid(1–40) vasoactivity by functional magnetic resonance imaging in mice. *Neuroscience* 155, 263–269.
- Luo, F., Wu, G., Li, Z., Li, S.J., 2003. Characterization of effects of mean arterial blood pressure induced by cocaine and cocaine methiodide on BOLD signals in rat brain. *Magn. Reson. Med.* 49, 264–270.
- Luo, Z., Yuan, Z., Tully, M., Pan, Y., Du, C., 2009. Quantification of cocaine-induced cortical blood flow changes using laser speckle contrast imaging and Doppler optical coherence tomography. *Appl. Opt.* 48, D247–255.
- Lyons, D., Friedman, D.P., Nader, M.A., Porrino, L.J., 1996. Cocaine alters cerebral metabolism within the ventral striatum and limbic cortex of monkeys. *J. Neurosci.* 16, 1230–1238.
- Mandeville, J.B., Ekstrom, L., Marota, J.J.A., Jenkins, B.G., Vanduffel, W., 2005. Contrast-enhanced fMRI of cocaine action in awake, non-human primate. *Proceedings of the International Society for Magnetic Resonance in Medicine*.
- Mandeville, J.B., Jenkins, B.G., Chen, Y.C., Choi, J.K., Kim, Y.R., Belen, D., Liu, C., Kosofsky, B.E., Marota, J.J., 2004. Exogenous contrast agent improves sensitivity of gradient-echo functional magnetic resonance imaging at 9.4T. *Magn. Reson. Med.* 52, 1272–1281.
- Mandeville, J.B., Marota, J.J., Kosofsky, B.E., Keltner, J.R., Weissleder, R., Rosen, B.R., Weisskoff, R.M., 1998. Dynamic functional imaging of relative cerebral blood volume during rat forepaw stimulation. *Magn. Reson. Med.* 39, 615–624.
- Marota, J.J., Mandeville, J.B., Weisskoff, R.M., Moskowitz, M.A., Rosen, B.R., Kosofsky, B.E., 2000. Cocaine activation discriminates dopaminergic projections by temporal response: an fMRI study in Rat. *Neuroimage* 11, 13–23.
- Mateo, Y., Budygin, E.A., John, C.E., Jones, S.R., 2004. Role of serotonin in cocaine effects in mice with reduced dopamine transporter function. *Proc. Natl. Acad. Sci. USA* 101, 372–377.
- Mueggler, T., Baumann, D., Rausch, M., Staufenbiel, M., Rudin, M., 2003. Age-dependent impairment of somatosensory response in the amyloid precursor protein 23 transgenic mouse model of Alzheimer's disease. *J. Neurosci.* 23, 8231–8236.
- Peoples, L.L., Gee, F., Bibi, R., West, M.O., 1998. Phasic firing time locked to cocaine self-infusion and locomotion: dissociable firing patterns of single nucleus accumbens neurons in the rat. *J. Neurosci.* 18, 7588–7598.
- Perles-Barbacaru, A.T., Procissi, D., Demyanenko, A.V., Jacobs, R.E., 2010. Nonlinear model for preprocessing of cerebral blood volume weighted functional MRI data and for evaluating pharmacokinetic properties of USPIO. *Proc. Intl. Soc. Mag. Reson. Med. 18th Annual Scientific Meeting, Stockholm (Sweden)*, p. 3550.
- Rocha, B.A., Fumagalli, F., Gainetdinov, R.R., Jones, S.R., Ator, R., Giros, B., Miller, G.W., Caron, M.G., 1998. Cocaine self-administration in dopamine-transporter knockout mice. *Nat. Neurosci.* 1, 132–137.
- Sanchez-Pernaute, R., Jenkins, B.G., Choi, J.K., Iris Chen, Y.C., Isacson, O., 2007. In vivo evidence of D3 dopamine receptor sensitization in parkinsonian primates and rodents with l-DOPA-induced dyskinesias. *Neurobiol. Dis.* 27, 220–227.
- Schmidt, K.F., Febo, M., Shen, Q., Luo, F., Sicard, K.M., Ferris, C.F., Stein, E.A., Duong, T.Q., 2006. Hemodynamic and metabolic changes induced by cocaine in anesthetized rat observed with multimodal functional MRI. *Psychopharmacol. Berl* 185, 479–486.
- Schwarz, A.J., Gozzi, A., Reese, T., Bifone, A., 2007a. In vivo mapping of functional connectivity in neurotransmitter systems using pharmacological MRI. *Neuroimage* 34, 1627–1636.
- Schwarz, A.J., Gozzi, A., Reese, T., Heidbreder, C.A., Bifone, A., 2007b. Pharmacological modulation of functional connectivity: the correlation structure underlying the pHfMRI response to d-amphetamine modified by selective dopamine D3 receptor antagonist SB277011A. *Magn. Reson. Imaging* 25, 811–820.
- Schwarz, A.J., Reese, T., Gozzi, A., Bifone, A., 2003. Functional MRI using intravenous contrast agents: detrending of the relative cerebrovascular (rCBV) time course. *Magn. Reson. Imaging* 21, 1191–1200.
- Schwarz, A.J., Zocchi, A., Reese, T., Gozzi, A., Garzotti, M., Varnier, G., Curcuruto, O., Sartori, I., Girlanda, E., Biscaro, B., Crestan, V., Bertani, S., Heidbreder, C., Bifone, A., 2004. Concurrent pharmacological MRI and in situ microdialysis of cocaine reveal

- complex relationship between the central hemodynamic response and local dopamine concentration. *Neuroimage* 23, 296–304.
- Shen, H.W., Hagino, Y., Kobayashi, H., Shinohara-Tanaka, K., Ikeda, K., Yamamoto, H., Yamamoto, T., Lesch, K.P., Murphy, D.L., Hall, F.S., Uhl, G.R., Sora, I., 2004. Regional differences in extracellular dopamine and serotonin assessed by in vivo microdialysis in mice lacking dopamine and/or serotonin transporters. *Neuropsychopharmacology* 29, 1790–1799.
- Sigovan, M., Bousset, L., Sulaiman, A., Sappey-Mariniere, D., Alsaïd, H., Desbleds-Mansard, C., Ibarrola, D., Gamondes, D., Corot, C., Lancelot, E., Raynaud, J.S., Vives, V., Laclede, C., Violas, X., Douek, P.C., Canet-Soulas, E., 2009. Rapid-clearance iron nanoparticles for inflammation imaging of atherosclerotic plaque: initial experience in animal model. *Radiology* 252, 401–409.
- Sora, I., Hall, F.S., Andrews, A.M., Itokawa, M., Li, X.F., Wei, H.B., Wichems, C., Lesch, K.P., Murphy, D.L., Uhl, G.R., 2001. Molecular mechanisms of cocaine reward: combined dopamine and serotonin transporter knockouts eliminate cocaine place preference. *Proc. Natl Acad. Sci. USA* 98, 5300–5305.
- Sora, I., Wichems, C., Takahashi, N., Li, X.F., Zeng, Z., Revay, R., Lesch, K.P., Murphy, D.L., Uhl, G.R., 1998. Cocaine reward models: conditioned place preference can be established in dopamine- and in serotonin-transporter knockout mice. *Proc. Natl Acad. Sci. USA* 95, 7699–7704.
- Thanos, P.K., Michaelides, M., Benveniste, H., Wang, G.J., Volkow, N.D., 2008. The effects of cocaine on regional brain glucose metabolism is attenuated in dopamine transporter knockout mice. *Synapse* 62, 319–324.
- Thomas, M.J., Kalivas, P.W., Shaham, Y., 2008. Neuroplasticity in the mesolimbic dopamine system and cocaine addiction. *Br. J. Pharmacol.* 154, 327–342.
- Thomsen, M., Hall, F.S., Uhl, G.R., Caine, S.B., 2009. Dramatically decreased cocaine self-administration in dopamine but not serotonin transporter knock-out mice. *J. Neurosci.* 29, 1087–1092.
- Uhl, G.R., Hall, F.S., Sora, I., 2002. Cocaine, reward, movement and monoamine transporters. *Mol. Psychiatry* 7, 21–26.
- Weissleder, R., Elizondo, G., Wittenberg, J., Rabito, C.A., Bengel, H.H., Josephson, L., 1990. Ultrasmall superparamagnetic iron oxide: characterization of a new class of contrast agents for MR imaging. *Radiology* 175, 489–493.
- Woods, R.P., Grafton, S.T., Holmes, C.J., Cherry, S.R., Mazziotta, J.C., 1998. Automated image registration: I. General methods and intrasubject, intramodality validation. *J. Comput. Assist. Tomogr.* 22, 139–152.
- Wu, E.X., Tang, H., Asai, T., Yan, S.D., 2004. Regional cerebral blood volume reduction in transgenic mutant APP (V717F, K670N/M671L) mice. *Neurosci. Lett.* 365, 223–227.
- Zocchi, A., Conti, G., Orzi, F., 2001. Differential effects of cocaine on local cerebral glucose utilization in the mouse and in the rat. *Neurosci. Lett.* 306, 177–180.